

Objective: To evaluate the effects of AT on cognition and regional brain glucose metabolism (rBGM) in MCI patients.

Methods: Subjects performed a twice a week, moderate intensity, AT program for 24 weeks. Assessments with ADAS-cog, a comprehensive neuropsychological battery, and evaluation of rBGM with positron emission tomography with ^{18}F -fluorodeoxyglucose (^{18}F FDG-PET) were performed before and after the intervention. Aerobic capacity was compared using the maximal oxygen consumption VO_2 max (mL/kg/min). ^{18}F FDG-PET data were analyzed on a voxel-by-voxel basis with SPM8 software.

Results: Forty subjects were included, with a mean (M) age of 70.3 (5.4) years and an initial Mini-Mental State Exam score of 27.4 (1.7). Comparisons using paired t-tests revealed improvements in the ADAS-cog (M difference: -2.7 (3.7), $p < 0.001$) and VO_2 max scores (M difference: 1.8 (2.0) mL/kg/min, $p < 0.001$). Brain metabolic analysis revealed a bilateral decrease in the rBGM of the dorsal anterior cingulate cortex (ACC), $p\text{FWE} = 0.04$. This rBGM decrease was negatively correlated with improvement in a visuospatial function/attentional test ($\rho = -0.31$, $p = 0.04$). Several other brain areas also showed increases or decreases in rBGM. Of note, there was an increase in the retrosplenial cortex, an important node of the default mode network, that was negatively correlated with the metabolic decrease in the dorsal ACC ($r = -0.51$, $p = 0.001$).

Conclusion: AT improved cognition and changed rBGM in areas related to cognition in subjects with MCI.

doi:10.1016/j.jns.2015.08.192

117

WFN15-0369

Mixed Topics 3

On the medical and neuropsychiatric phenomena depicted at two famous medieval marian miracle spanish books

F.A.A. Gondim^a, W. Gondim^b. ^aInternal Medicine, Universidade Federal do Ceará, Fortaleza, Brazil; ^bSaúde Materna e Infantil, Universidade Federal do Ceará, Fortaleza, Brazil

Background: The XIII Century is known for the highest peak of devotion to Holy Mary (Marian Century). We have recently reviewed the neuropsychiatric aspects depicted at the medieval Spanish book “Cantigas de Santa Maria” (Neurology 2015;84:1991–6).

Objective: To report and compare the medical and neuropsychiatric findings in 2 famous medieval books of Marian miracles.

Material and methods: We reviewed all the miracles depicted at the XIII Century books “Cantigas de Santa Maria” (CSM, written in Galician at the court of Alfonso X) and Milagros de Nuestra Señora (MNS), written by Gonzalo de Berceo (ca. 1197–1264), the first Castilian poet known by name and compared the relevant medical, neurological and neuropsychiatric events in both books.

Results: MNS does not have illuminations and at least 9 similar narratives were also found on CSM. Among the 25 miracles reported at MNS, 36% (9/25) included medically relevant facts with 4 additional examples of resurrection and 2 stories describing evil possession. The most common medical/neuropsychiatric subjects included ob/gyn subjects ($N = 2$), sudden death (MI?), intellectual decline, alcohol intoxication, suicide, infanticide, infections and uncorrupted body after death ($N = 1$ for each). At the 427 canticles from CSM (353 miracles), 270 medically relevant facts (187 canticles) were found. Possible neuropsychiatric conditions were described in 98 canticles. Blindness and dystonia/weakness/deformities were the most common neurological phenomena. Several examples were also detailed by illuminations.

Conclusion: Medically relevant facts were described in both books. Accounts of neuropsychiatric disorders were more complete on CSM

(including illuminations) whether examples on MNS were mostly psychiatric.

doi:10.1016/j.jns.2015.08.193

118

WFN15-0199

Mixed Topics 3

Chasing dizzy chimera: Diagnosis of combined peripheral and central vestibulopathy

J. Kim^a, S. Choi^b, H. Kim^c. ^aNeurology, Seoul National University, College of Medicine, Seongnam, South Korea; ^bNeurology, Eulji University Hospital, Daejeon, South Korea; ^cBiomedical Laboratory Science, Kyungdong University, Goseong, South Korea

Background and objectives: Diagnosis of combined peripheral and central vestibulopathy remains a challenge since the findings from peripheral vestibular involvements may overshadow those from central vestibulopathies or vice versa. The aim of this study was to enhance the detection of these intriguing disorders by characterizing the clinical features and underlying etiologies.

Methods: We had recruited 55 patients with combined peripheral and central vestibulopathy at the Dizziness Clinic of Seoul National University Bundang Hospital from 2003 to 2013. Peripheral vestibular involvement was determined by decreased caloric responses in one or both ears, and central vestibulopathy was diagnosed with obvious central vestibular signs or the lesions documented on MRIs to involve the central vestibular structures.

Results: Combined peripheral and central vestibulopathy could be classified into four types according to the patterns of vestibular presentation. Infarctions in the territory of anterior inferior cerebellar artery were the most common cause of acute unilateral cases while cerebellopontine angle tumors were mostly found in chronic unilateral ones. Wernicke encephalopathy and degenerative disorders were common in acute and chronic bilateral disorders. HINTS may not detect central lesions in combined vestibulopathy. The dissociation in the abnormalities between the caloric and head impulse tests may suggest a combined peripheral and central vestibulopathy.

Conclusions: Peripheral vestibular signs may overshadow the central ones in combined peripheral and central vestibulopathy. Given the requirements for urgent treatments and potentially grave prognosis of combined vestibulopathy, central signs should be sought carefully even in patients with obvious clinical or laboratory features of peripheral vestibulopathy.

doi:10.1016/j.jns.2015.08.194

119

WFN15-1206

Mixed Topics 3

Recurrent benign paroxysmal positional vertigo: Analysis of 170 patients

F. Lerchundi, A.H. Laffue, M. Olivier, F.J. Gualtieri. Neurology, FLENI, Buenos Aires, Argentina

Background: Benign paroxysmal positional vertigo (BPPV) accounts for 20% of the diagnosis in specialized centers. The recurrence after an effective treatment has been reported in 20–50%.

Objective: To assess the rate of recurrence of BPPV and search of related comorbidities.

Patients and methods: Retrospective chart review of 170 patients with recurrent BPPV, analysis of epidemiological characteristics and

the number of canalith repositioning procedures (CRP). The definition of recurrence was any new attack that occurred at least one month after the resolution of the first BPPV. The chi-square test was used to assess the relationship between comorbidities and recurrence BPPV. Statistical significance was defined as $p < 0,05$.

Results: 674 patients had diagnosis of BPPV between 2007 and 2011 in our neurology unit, 170 (25%) were defined as recurrent BPPV. The mean age was 62 years and 125 (73.5%) were female. Posterior semicircular canal was involved in 89% of recurrences. The media of recurrence was 2.04 BPPV episodes per patient. The number of CRP increased along with patient's age. Regarding the evaluated comorbidities: 43% had arterial hypertension, 37.5% high levels of blood lipids, 30% migraine, 24.2% stroke, 21.2% hypothyroidism, 22% tensional headache, 18.8% head trauma, 14.7% diabetes and 11% smoked. A value of chi squared of 8.12 was calculated for lacunar stroke in patients with recurrent BPPV ($p = 0.004$) and for patients with stroke, was 6.88, $p = 0.009$.

Conclusion: A relationship between microangiopathic damage and BPPV recurrence has been observed, but more studies are needed in order to demonstrate the validity of this hypothesis.

doi:10.1016/j.jns.2015.08.195

120

WFN15-0275

Mixed Topics 3

Microperimetry in the diagnosis of the first manifestation of optic neuritis in multiple sclerosis

E. Ioyleva, M. Krivosheeva. Neuroophthalmology, S. Fyodorov Eye Microsurgery, Federal State Institution, Ministry of Health, Moscow, Russia

Background: We patented a new method of diagnosis of optic neuritis by microperimetry (Patent RU No. 2548511, priority of 29.11.2013).

Objective: To identify subclinical lesions of the optic nerve at the first manifestation multiple sclerosis.

Methods: The clinic S. Fyodorov Eye Microsurgery, Moscow, examined seven patients with complaints typical for monolateral optic neuritis (ON). RC-5000 Tomey, Cirrus HD-OCT; Humphrey HFA II-750i; Microperimetry on MP-1 (program 12° 10 dB stimulus Goldmann III), MRI of the brain and orbits (Siemens 1.5 Tl).

Results: Visual acuity 7 patients in both eyes 1.0, fundus – unaffected. Intact fellow eye. OCT: retina nerve fiber layer from 88 to 108 nm, and the volume of the retinal ganglion cells 60 to 73 nm. Microperimetry: 4 eyes – a significant decrease in the average sensitivity of 6, 2 to 10, 8 dB and paracentral defects within 0, 2 and 4° from the fixation target, and 3 eyes – a moderate decrease in the average sensitivity from 15.2 to 16.8 dB, paracentral defects within the 2, 4, 6°, preferably in the upper quadrant. MRI in all patients: periventricular, subcortical, multiple foci size of 0.3–0.7 cm, some

with signs of perifocal edema in the form of activity. Neurologist diagnosed with multiple sclerosis was confirmed.

Conclusions: Method microperimetry revealed paracentral defects in cases of subclinical lesions of the optic nerve. In all cases had been diagnosed with multiple sclerosis. Microperimetry – informative method for diagnosis of subclinical lesions of the optic nerve at the first manifestation of multiple sclerosis.

doi:10.1016/j.jns.2015.08.196

121

WFN15-0911

Mixed Topics 3

Clinical and neuro-ophthalmologic predictors of visual outcome in idiopathic intracranial hypertension (IIH)

A. Kapila^a, V. Lal^a, A. Gupta^b. ^aNeurology, PGIMER, Chandigarh, India; ^bOphthalmology, PGIMER, Chandigarh, India

Background: Once thought benign, IIH is now recognized as a potentially vision devastating entity. Predictors of worse outcome may aid in appropriate management of patients of IIH.

Objective: To assess visual morbidity in patients of IIH in terms of clinical and neuro-ophthalmologic parameters.

Methods: In a prospective study of 40 patients of IIH, clinical and neuro-ophthalmologic parameters were noted. All parameters were graded using objective criteria and compared with final visual outcome of patients to determine any correlation.

Results: Presence of headache and transient visual obscurations (TVOs) at presentation was significantly associated with good visual outcome. Higher CSF opening pressure, Increased Retinal nerve fibre layer (RNFL) thickness ($> 178 \mu\text{m}$) on optical coherence tomography, abnormal contrast sensitivity at high frequency and prolonged P100 latency on visual evoked potential (VEP) were associated with poor visual outcome. No definite correlation between presence of cranial nerve palsies, other clinical features, obesity, grade of papilledema and findings on neuroimaging with final visual outcome was established.

Conclusion: Visual Outcome in patients of IIH can be predicted at presentation by various parameters providing window for early and adequate treatment. High Body mass index, high CSF opening pressure, greater RNFL thickness, abnormal VEP or contrast sensitivity are some of the alarming signs for physicians necessitating need for aggressive monitoring and management. On the contrary, presence of headache or TVOs at presentation may predict a better visual outcome. Close follow up of the patients with respect to the neuro-ophthalmologic parameters as described may help in segregating the potential candidates for aggressive management early in the disease course.

doi:10.1016/j.jns.2015.08.197