

Palermo and AOUP Palermo, Palermo, Italy; ^bRobotics Lab Dept of Chemical Management Informatics and Mechanical Engineering, University of Palermo, Palermo, Italy; ^cALS Clinical Research Center, University of Palermo, Palermo, Italy

Background. Amyotrophic lateral sclerosis (ALS) leads to a complete limb paralysis, dysphagia and anarthria. A brain-computer interface (BCI) technology may aid ALS patients in communication and motor control.

Objective. To set up a BCI system for wireless control, by ALS patients, of a humanoid robot, with the aim to reach and grasp a glass of water.

Patients and methods. Four non-demented ALS patients were recruited. Controls were four healthy subjects, matched for demographic variables. A BCI command interpreter was used to control a humanoid robot. The task was to instruct the robot to move towards a glass of water, reach and grasp it (first item) and then bring the glass to the subject (second item). The protocol consisted of a calibration session, an online session, in which the two items are sequentially selected, and a robot session where the two items translate into high level commands. The minimal accuracy of each response and the number of errors each session were evaluated and analysed.

Results. All ALS patients completed the task (5 trials, 95.5% success). Controls performed comparably with a 100% success over the 5 trials. The minimum accuracy leading to a correct item selection for the robot movement was slightly better for ALS patients (ALS 60% vs controls 53%, $p = 0.49$).

Conclusions. ALS patients can successfully control a humanoid robot through a BCI system. This bears the potential to virtually restore the autonomous motion of an ALS patient, enabling him to extend his presence beyond the boundaries of his bed.

doi:10.1016/j.jns.2015.08.200

125

WFN15-1342

Motor Neuron Disease 1

Fibroblasts from patients with amyotrophic lateral sclerosis (ALS) associated with mutations in *tardbp* gene as model of TDP-43 proteinopathy

C. Hewamadduma^{*}, A. Grierson, A. Higginbottom, P. Shaw. *Academic Neurology, Sheffield Institute for Translational Neurosciences, Sheffield, United Kingdom*

Rationale & hypothesis: ALS is a devastating progressive neurodegenerative condition, which results in death. TARDBP encoded protein, TDP-43, has been implicated in both the sporadic and familial ALS cases. Animal models of TDP-43 have been inconclusive and the role of TDP-43 in ALS remains an enigma to date. Fibroblasts obtained from the patients carrying mutations in TARDBP gene provide a vital tool in investigation of TDP-43 due to physiological levels of TDP-43.

Methodology: Immunocytochemistry was performed on three lines of control and three different TDP-43 mutant fibroblast lines (M337V, G287V, A321V) and confocal microscopy was performed to identify general TDP-43, phosphorylated TDP-43 and anti p62 (to identify ubiquitin) localisations. Fibroblasts were also subjected to 0.5 mM arsenite and the stress response was assessed using markers of stress granules such as TIAR and HUR. Recovery after stress was also assessed.

Findings/conclusion: In keeping with findings in ALS postmortem material, relative clearing of nuclear TDP-43 was noted in mutant TDP-43 fibroblasts ($p < 0.001$). TDP-43 fibroblasts also showed accumulation of p62 positive aggregates ($p < 0.0003$), and phosphorylated TDP-43 accumulation ($p < 0.001$) compared to controls, suggesting that mutant

TDP-43 fibroblasts share some characteristics of the surviving motor neurons from both sALS and fALS. Following exogenic stress endogenous TDP-43 localised to HUR positive stress granules ($p < 0.01$), formation of stress granules and their recovery were significantly impaired in mTDP-43 cases ($p < 0.01$) suggesting that dysfunction of TDP-43 dysregulates handling of exogenic stress. We suggest that this may contribute to premature degeneration of motor neurons expressing mutant TDP-43 in ALS patients. Fibroblasts also form a robust and an economical platform to study TDP-43 related neurodegeneration.

I have obtained patient and Institutional Review Board (IRB) approval and local ethics committee approval.

doi:10.1016/j.jns.2015.08.201

126

WFN15-1481

Motor Neuron Disease 1

C9ORF72 repeated expansion in patients with familial amyotrophic lateral sclerosis from a Brazilian research center. A preliminary report

G. Chadi^{*}, J.R. Maximino, F.M.H. Jorge, D. Callegaro. *Neurology, University of Sao Paulo School of Medicine, São Paulo, Brazil*

Background: The expanded GGGGCC hexanucleotide repeat in the non-coding region of the chromosome 9 open reading frame 72 (*C9ORF72*) gene is the most common genetic abnormality in familial amyotrophic lateral sclerosis (FALS).

Objective: To determine the *C9ORF72* hexanucleotide repeat expansion in FALS patients from ALS Unit of São Paulo, Brazil.

Patients and methods: Patients with FALS from the ALS Unit of Clinics Hospital, University of São Paulo Medical School, Brazil have been evaluated for the presence of an expanded (GGGGCC) in *C9ORF72*. A repeat-primed-PCR reaction was applied to provide a qualitative assessment of the expansions. PCR products were analyzed on an ABI3730 and visualized using GeneMapper-software. A cutoff of >30 repeats combined with a typical sawtooth pattern was considered pathologic.

Results: Preliminary results from 15 FALS patients (mean age of onset 51.40 ± 3.02 years) are shown. The repeat expansion was present in 5 FALS cases (33.3%) and 10 FALS did not present the pathologic expansion. One patient with *C9ORF72* expansion presented a bulbar-onset and developed later a frontotemporal degeneration. Patients without *C9ORF72* expansion had a spinal-onset disease. FALS patients with *C9ORF72* expansion developed a later onset symptoms (54.80 ± 4.16 years) when compared to FALS without expansion (49.7 ± 4.07 years). A shorter lifespan was seen in *C9ORF72* expansion carriers (5.0 ± 1.22 years) than *C9ORF72* negative (9.10 ± 2.42 years).

Conclusion: A high frequency of *C9ORF72* expansion was detected in this partial report of a small FALS sample of São Paulo ALS Unit. Supported by FAPESP and CNPq, Brazil.

doi:10.1016/j.jns.2015.08.202

127

WFN15-0089

Motor Neuron Disease 1

Epidemiological and clinical features of amyotrophic lateral sclerosis in Uzbekistan

D. Mirzaeva^{*}, A. Prohorova, H. Daminova. *Neurology disease, Tashkent Medical Academy, Tashkent, Uzbekistan*

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease characterized by multifactorial etiology, affections of central and

peripheral motor neurons, progressive course and onset of terminal respiratory insufficiency. The amount of patients including ones in Uzbekistan that are suffering and dying from ALS is increasing. The development of a familial ALS (FALS) shows the role of mutations of the gene producing the superoxide dismutase 1.

Data on morbidity of ALS on base of register for the period of 2013–2014 was studied. Special attention was paid to geographic epidemiology of ALS in different regions of Uzbekistan and to features of clinical manifestations of this disease in Uzbek nationality.

During the analyzed period there were 3.4 cases of ALS per 100,000 per year registered. 70% were patients from Andijan valley, 30% from other regions. Gender analysis determined that 60% of patients were women (mean age 43 ± 4.2 years), 40% men (mean age 48 ± 3.4 years). Most patients associated the debut of ALS with virus disease. Disease of 80% of patients debuted by bulbar form, of 20% by spinal form. The FALS is absent in our country. The average life expectancy was 2.5 ± 0.45 years during the bulbar form and 3 ± 0.86 years during the spinal form. The onset of decompensation period on the average 6 month later after debut.

The analysis of the register of ALS in Uzbekistan demonstrated that ALS was the most common disease in Andijan valley, primary occurred in women, is not of a personal nature and mainly declares itself by bulbar form.

doi:10.1016/j.jns.2015.08.203

128

WFN15-0254

Motor Neuron Disease 1

Rehabilitation of blood-spinal cord-barrier toward amyotrophic lateral sclerosis therapy

B. Solomon^{*}, I. Rabinovich-Nikitin. *Biotechnology-Microbiology, Tel Aviv University, Tel-Aviv, Israel*

Background: ALS patients and transgenic mice expressing ALS-associated superoxide dismutase 1 (SOD1) mutations show alterations in the blood-spinal cord barrier (B-SC-B) as suggested from the reduction of levels of various tight junction proteins (TJPs) including zonula occludens-1 (ZO-1), occludin and claudin-5 between endothelial cells and early protection of the B-SC-B integrity was found to delay onset of motor-neuron impairment and degeneration.

Objective: The aim of this research was to investigate if inhibition of the axis CXCL12/CXCR4 receptor widely expressed in neurons and glial cells and modulates neuronal apoptosis, may improve motor neurons survival by increasing the expression of tight junction proteins and rehabilitation of the barrier.

Materials and methods: Transgenic mice model of ALS were treated with AMD3100, antagonist of CXCR4. Motor function, weight changes and survival were evaluated. In a separate experiment, mice were sacrificed after one month of treatment and levels of proteins essential for the formation of the barrier in comparison with proteins that do not participate in the barrier were measured.

Results: We found that chronic administration of AMD3100 to ALS mouse model was effective in restoring the expression of tight

junction proteins and considerably increase the survival, confirming the importance of early treatment for rehabilitation of the barriers to prevent infiltration of neurotoxic products and microhemorrhages.

Conclusions: These data reveal that multi-faceted action of AMD3100 may provide a novel option for ALS therapy leading to rehabilitation of B-SC-B proteins and thus preventing additional damage to motor neurons.

doi:10.1016/j.jns.2015.08.204

129

WFN15-1264

Motor Neuron Disease 1

Far beyond our typical dengue fever on three cases reported: weakness, visual loss and aphasia as initial clinical presentations?

A. Cronemberger-Andrade^{a,*}, H.R. Soares-Neto^a, A.F.P. Pouza^a, D.D. de-Faria^a, L. Dongyang^a, N.C. Bergamasco^a, C.A. de-Albuquerque^b, E.P. de-Andrade^c, S.M.C.A. Silva^a. ^aServiço de Neurologia, Hospital do Servidor Público Estadual Francisco Morato de Oliveira (HSPE-FMO/IAMSPE), São Paulo, Brazil; ^bServiço de Oftalmologia, Hospital do Servidor Público Estadual Francisco Morato de Oliveira (HSPE-FMO/IAMSPE), São Paulo, Brazil; ^cServiço de Neuro-oftalmologia, Hospital do Servidor Público Estadual Francisco Morato de Oliveira (HSPE-FMO/IAMSPE), São Paulo, Brazil

Background. The mosquito-borne dengue virus human infection is endemic in Brazil and the most populated São Paulo State concentrates a great number of cases. There is a wide range of possible clinical manifestations. Secondary dengue-related diseases and complications can show up with nonspecific symptoms to more severe hemorrhagic shock. Possible neurological manifestations are also a part of initial clinical assessment of an infected patient and can define severity of affection. Objective. This cases report aims to study complications dengue fever can present with and alert health care professionals dealing with this disease.

Patients. Here we present three cases of adult patients admitted in March 2015 at our hospital during a current epidemic of dengue viral infection in the State who experienced neurological complications in the convalescent phase of infection: a man with weakness (Guillain-Barré syndrome), a woman with visual loss (unilateral maculopathy), and a woman with aphasia (post-seizure Todd's palsy). The patients had in common a previously acute febrile disease diagnosed as dengue fever.

Results. After specific investigation and treatment, all of them had their complaints improved.

Conclusion. These various disease presentations on patients referred to neurological care should be promptly diagnosed and treated. To call attention and inform health teams on these different presentations is essential. The dissemination of new clinical guidelines for health professionals on non-tertiary services including these neurological complications might be useful and can yield better outcomes.

doi:10.1016/j.jns.2015.08.205