

peripheral blood. Meanwhile, relevant cytokines were isolated in peripheral blood to evaluate functions of Treg and Th17, respectively.

Results: Cerebral hemorrhage occurs in half of patients as an onset symptom, followed by cerebral ischemia. Our data revealed that both percentage of Treg and Th17 among lymphocytes was elevated in MMD patients. Similarly, MMD patients showed significant increase in functions of Treg and Th17 as evidenced by increased expression of IL-6, IL-10, IL-12, TNF- α , VEGF and TGF- β in peripheral blood. However, no significant difference in the balance of Treg/Th17 was detected between two groups. Furthermore, onset symptoms and gender were independent with these changes.

Conclusion: We first report here increased expression in Treg and Th17 cells that were found in MMD patients, which may provide valuable insight into the immune-related pathology of MMD.

doi:10.1016/j.jns.2015.08.326

251

WFN15-0944

Stroke 1

Education of children about stroke and heart attack: Feasibility and effectiveness pilot study

P. Sobotkova^a, V. Svobodova^a, H. Pokorna^a, O. Suchy^b, M. Seifert^c, B. Zuchova^d, J. Neumann^e, J. Vitovec^f, R. Mikulik^g. ^aInternational Clinical Research Center, St. Anne's University Hospital in Brno, Brno, Czech Republic; ^bNational Institute for Education, National Institute for Education Prague Czech Republic, Prague, Czech Republic; ^cInternational Clinical Research Center, St. Anne's University Hospital in Brno, Brno, Czech Republic; ^dEmergency Medical Services, Emergency Medical Services of South Moravian, Brno, Czech Republic; ^eDepartment of Neurology, County Hospital Chomutov, Chomutov, Czech Republic; ^fDepartment of Cardiology, St. Anne's University Hospital in Brno, Brno, Czech Republic; ^gDepartment of Neurology and International Clinical Research Center, St. Anne's University Hospital in Brno, Brno, Czech Republic

Background and objective: The effectiveness of stroke educational campaigns aimed at adults is limited. The study objective was to evaluate the feasibility and efficacy of an innovative stroke and heart attack educational program for children.

Methods: The 45 minute multimedia web-based videoprogram was designed to teach the correct response to stroke and heart attack symptoms and to test short-term and long-term retention. Population consisted of school children aged 13–15 from 2 remote counties (target and control) in the Czech Republic. Target population obtained education and testing. Control population had only testing. Ethical committee approved the protocol and children's parents signed informed consent.

Results: All children aged 13–15 participated from 37 schools (8% of all basic schools in the county) as target population (n = 2436) during 2014/2015. The control group involved 426 pupils from 6 schools. The baseline knowledge (measured as % of correct response to 12 questions/simulation video-clips) was the same in the target and control population (59% versus 58%). After education, knowledge in the target population improved (67%, paired p = 0.001). After 3 months, knowledge was 61% in the target and 53% in the control populations (intergroup difference p = 0.001). Knowledge about heart attack was higher than for stroke: at baseline in both target (79% versus 57%) and controls (74% versus 58%) as well as at 3 months (77% versus 57%).

Conclusion: School education on stroke is feasible. Knowledge about stroke is worse than for heart attack. The educational program increased knowledge and response to stroke and heart attack symptoms in the short-term and also long-term.

doi:10.1016/j.jns.2015.08.327

252

WFN15-1006

Stroke 1

A national survey of preferences for mobile applications (APPS) among stroke survivors & caregivers

N. Gilles^a, S. Zelonis^a, L. Beving^b, D. Burton^c, C. Ventura-DiPersia^c, S. Kunnakkat^d, C. Balucani^d, M. White^e, R. Browne^e, S. Levine^a. ^aNeurology Stroke Center, SUNY Downstate Medical Center, Brooklyn, USA; ^bStroke, National Stroke Association, Centennial, CO, USA; ^cPublic Health, Hunter College, CUNY, NY, USA; ^dNeurology/Stroke Center, SUNY Downstate Medical Center, Brooklyn, USA; ^eResearch, Arthur Ashe Institute for Urban Health, Brooklyn, USA

Background: Mobile technology is underutilized in patient-centered healthcare. Mobile applications (apps) may allow stroke survivors/caregivers to actively participate in stroke-related care/needs.

Objective: Investigate preferences of stroke survivors and caregivers for mobile apps to facilitate post-stroke care/needs.

Subjects & methods: Nationwide survey of 17 questions distributed to 11,720 stroke survivors/caregivers identified from National Stroke Association's database via e-/postal mail, including introduction letter and IRB-approved consent. The survey was developed using formative focus groups in Brooklyn and Colorado. Qualitative information about current smartphone usage/interest in health-related apps was collected. Preferences were explored by gender/race/ethnicity/age/education using χ^2 .

Results: 1221 survivors and 396 caregivers responded (14% response). Scheduling doctor/rehab appointments was preferred by survivors and caregivers of all ethnicities, with doctor appointments/medication reminder/blood pressure tracking as most favored by both. App usefulness declined with age ($\chi^2 = 19.7$, p = 0.02). App with rehab exercises was more cited by younger survivors with older wanting trustworthy medical information ($\chi^2 = 41.4$, p < 0.001). App to find local stroke-related resources was preferred by the majority of caregivers and stroke survivors. More Afro-Caribbean/American survivors (42%) preferred use of stroke support groups vs. Hispanics (36%) or Caucasians (28%), $\chi^2 = 45.1$, p < 0.001 as did younger ($\chi^2 = 41.3$, p < 0.001). App tracking fitness/diet was more desired by the majority of stroke survivors than caregivers. App facilitating stroke survivors' communication was highly favored by survivors while not considered useful by the majority of caregivers.

Conclusions: Developing relevant apps requires feedback from users. We identified useful key features reported by stroke survivors and caregivers to build a stroke-dedicated app.

doi:10.1016/j.jns.2015.08.328

253

WFN15-1024

Stroke 1

Mobile devices for remote acute stroke neuroimaging interpretation: Diagnostic accuracy

C. Balucani^a, Q. Hao^a, J. Weedon^b, S. Levine^a, R. Study Group^c. ^aNeurology Stroke Center, SUNY Downstate Medical Center, Brooklyn, USA; ^bBiocomputing & Statistics, SUNY Downstate Medical Center, Brooklyn, USA; ^cNeurology & Radiology, The REliability of MOBILE Technologies for Acute Stroke Neuroimaging Data Interpretation (REMOTE) Study Group, Brooklyn, USA

Background: Diagnostic accuracy of various mobile devices for remote acute ischemic stroke CT scan interpretation is not known.

Objective: Compare diagnostic accuracy of different mobile devices in acute CT interpretation for various ischemic stroke findings among various readers.

Subjects/methods: After IRB approval, 20 selected AIS CTs were independently interpreted by 11 readers (7 neurologists/3 neuroradiologists/1 radiologist) on 2 mobile devices: iPad®2 (1024 × 768 pixels) and iPhone®4 (960 × 640 pixels), and on a Picture Archiving and Communication System (PACS) workstation (1280 × 1024 pixels). Mobile devices used FDA-cleared ResolutionMD software (ResMD® Calgary Scientific, Calgary, Canada). Primary outcome measures: any acute ischemic sign (AIS), any non-acute ischemic sign (NAIS), and hyperdense middle cerebral artery (HMCA) sign. Intra-rater and inter-device accuracy, using a rater specific gold standard, was evaluated; estimated % sensitivity (Sen) and % specificity (Spe) with 95% Agrest-Coull confidence intervals (CIs) were calculated.

Results: Across all 11 readers, for AIS: Sen = 76 (68, 82) and Spe = 75 (64, 83) on iPhone; Sen = 76 (69, 83) and Spe = 71 (60, 80) on iPad. For NAIS: Sen = 80 (72,86) and Spe = 77 (67, 84) on iPhone; Sen = 75 (67, 82) and Spe = 80 (71, 87) on iPad. For HMCA: Sen = 53 (39, 67) and Spe = 97 (93, 99) on iPhone; Sen = 62 (48, 75) and Spe = 95 (91, 98) on iPad.

Conclusions: We believe this is the first study directly comparing the diagnostic accuracy of two different mobile devices for AIS on head CT. Both iPhone and iPad have fair sensitivity for detection of AIS and NAIS, but poorer sensitivity for HMCA when read by readers from different specialties. Specificity was good–excellent for all 3 CT findings. iPhone and iPad have similarly good diagnostic accuracy for AIS supporting their use for remote neuroimaging interpretation. Validation in clinical practice needs to be confirmed.

doi:10.1016/j.jns.2015.08.329

254

WFN15-1036

Stroke 1

Comparative analysis of recanalization treatments in basilar artery occlusion

P.J. Lindsberg^a, T.R. Sairanen^a, S. Nagel^b, H. Silvennoinen^c, O. Salonen^c, D. Strbian^a. ^aDepartment of Neurology, Helsinki University Hospital, Helsinki, Finland; ^bDepartment of Neurology, University Hospital Heidelberg, Heidelberg, Germany; ^cHelsinki Medical Imaging Center, Helsinki University Hospital, Helsinki, Finland

Background: Basilar artery occlusion (BAO) is the most devastating form of stroke, and the current wisdom is to reverse it with aggressive revascularization treatments. Recent trials using endovascular stentriever have shown efficacy in anterior circulation, but did not recruit posterior circulation occlusions. The preferred approach in BAO remains unknown.

Objective: This study collected published series of recanalization treatments in BAO to determine relative efficacy.

Patients and methods: We analyzed systematically the reported outcomes produced in 17 cohorts published from 2005 comprising 803 patients. Predictors of futile recanalization (FR; 3-month mRS score 4 to 6) were determined in the largest cohort (162, Helsinki).

Results: Superior results were achieved by primary thrombectomy with recanalization in 91% and good outcome (mRS 0 to 2) in 36%. Still, there was a substantial FR rate at 60%, which was improved using modern stentriever only (52.8%). Good outcome was reported by mechanical approaches either alone or on-demand more frequently

than by exclusively pharmacological protocols (35.5% vs. 24.4%, $p < 0.001$) in line with higher recanalization rates (84.1% vs. 70.9%, $p < 0.001$). In the largest single-center cohort, the most significant predictor was extensive baseline ischemia, increasing the odds of futility 20-fold (95% CI 4.4–92.3, $p < 0.001$). Ventilation support increased FR 7-fold (2.1–23.7, $p < 0.01$).

Conclusion: Modern stentriever approaches have reported superior outcome rates over exclusively pharmacological thrombolysis protocols in BAO. However, still more than half of the recanalizations turn out futile. To improve this, more careful patient selection is needed, especially to exclude patients with an already extended baseline ischemia.

doi:10.1016/j.jns.2015.08.330

255

WFN15-1080

Stroke 1

PIAST – Platelet inhibition assessment in stroke trial multiplate analyzer based assessment of the efficacy of antithrombotic medication in stroke patients

S. Menon^a, T. Jun^b, U. Meyding-Lamadé^c, A. Grabowski^d. ^aNeurology, Klinikum Aschaffenburg-Alzenau, Aschaffenburg, Germany; ^bInternal Medicine, Krankenhaus Nordwest GmbH, Frankfurt, Germany; ^cNeurology, Krankenhaus Nordwest GmbH, Frankfurt, Germany; ^dNeurology, Neuropraxis, Frankfurt, Germany

Objective: The aim of this prospective trial was to assess 1) the percentage of patients being treated with antiplatelet medications (AM) prior to or for secondary prevention of stroke who disclose an insufficient platelet aggregation (PA) inhibition assessed with Multiplate (MEA) – a validated platelet function test – and 2) conditions that might contribute to PA.

Background: A significant number of patients suffer from a recurrent cerebral ischemia despite treatment with AM without detectable progression of the underlying illness likely due to antiplatelet non-responsiveness caused by different factors. Platelet function was analyzed through MEA to identify the effects of antiplatelet therapy in stroke patients. Furthermore, the question if aspirin related insufficient platelet inhibition is correlated with the clinical diagnosis is addressed.

Methods: We analyzed the platelet function of 455 patients admitted with a suspected stroke either prior to being treated with aspirin and clopidogrel or being newly placed on AM employing MEA.

Results: Data shows that 120 (26.3%) of all examined patients with suspected ischemic stroke fall into the category of potential resistance to antiplatelet agents, mounting to 30% of the eventually confirmed 400 cerebral ischemia patients. Increased inflammatory biomarkers and use of NSAIDs was associated with an elevated PA and stroke/TIA incidence. Diabetes mellitus and decreased renal function do not seem to have an impact though. Insufficient PA seems to correlate with an increased rate of cerebral ischemia.

Conclusion: Nearly 1/3 of cerebral ischemia patients are potential aspirin nonresponders, which might have significant implications in the medical treatment of stroke patients.

I have obtained patient and/or Institutional Review Board (IRB) approval, as necessary

doi:10.1016/j.jns.2015.08.331