



Stroke 3

264
WFN15-0253
Stroke 3

Thrombus histology and cryptogenic stroke – a different approach to determine etiology

T. Boeckh-Behrens^a, J. Kleine^a, C. Zimmer^a, H. Poppert^b. ^aDepartment of Neuroradiology, Klinikum rechts der Isar, München, Germany; ^bDepartment of Neurology, Klinikum rechts der Isar, München, Germany

Objective: Ischemic stroke of undetermined cause is considered to be a major health issue and therefore recommended to be a top research focus in the field. Histopathological analysis of human “in-vivo”, mechanically extracted thrombi of patients with large vessel occlusion may contain additional “in-depth” information concerning the underlying pathology. Aim of this study was to elucidate the clinical problem of cryptogenic stroke by use of a multiparametric approach, analyzing histological, interventional and clinical outcome data of stroke patients.

Methods: Thrombi of 145 consecutive patients with large vessel occlusion were collected during mechanical recanalization. The HE-stained specimen was analyzed concerning overall appearance and relative quantitative component fractions of erythrocytes, fibrin/thrombocytes and leucocytes. Detailed clinical and interventional data were obtained. Statistical analysis was performed searching for significant differences or pattern similarities in different stroke subtypes, defined by the international TOAST criteria. IRB and patient approval has been obtained.

Results: In a consistent and conclusive overall pattern, arterio-embolic and TOAST 4 (mostly dissections) differ significantly from cardio-embolic and cryptogenic stroke subtype in histological parameters (e.g. lower erythrocyte fraction in cardio-embolic and cryptogenic groups, $p = 0.038$), interventional (e.g. higher number of maneuvers in cardio-embolic and cryptogenic groups, $p = 0.012$) and outcome (e.g. higher mRS in cardio-embolic and cryptogenic groups, $p = 0.011$) data. Cardio-embolic and cryptogenic subtype show matchable thrombus structure and similar clinical and interventional characteristics.

Interpretation: The findings of this multiparametric approach strongly support the hypothesis of predominantly cardio-embolic mechanisms accounting for the majority of cryptogenic strokes.

doi:10.1016/j.jns.2015.08.340

265
WFN15-0190
Stroke 3
Blood based biomarkers to identify subclinical brain damage in essential hypertension

A. González-Quevedo^a, S. González-García^a, O. Fernández Concepción^b, M. Peña-Sánchez^a, L. Quevedo-Sotolongo^c, R. Santiesteban-Freixas^d,

M.C. Menéndez-Saínez^e, R.A. Fernández-Carriera^e, I. Fernández-Almirall^e, Z.M. Hernández-Díaz^f. ^aDepartment of Neurobiology, Institute of Neurology and Neurosurgery, Havana, Cuba; ^bDepartment of Neurology, Institute of Neurology and Neurosurgery, Havana, Cuba; ^cDepartment of Imaginology, “Cira García” Central Clinic, Havana, Cuba; ^dDepartment of Neuroophthalmology, Institute of Neurology and Neurosurgery, Havana, Cuba; ^eDepartment of Neurobiology, Institute of Neurology and Neurosurgery, Havana, Cuba; ^fDepartment of Neuroimaging, Institute of Neurology and Neurosurgery, Havana, Cuba

Background: Neuroimaging has demonstrated that subclinical brain damage in essential hypertension is more prevalent than cardiovascular or renal impairment; nevertheless, screening for nervous system involvement is difficult due to low accessibility and high cost techniques.

Objective: To explore if hemochemical markers of brain damage and inflammation could predict the presence of subclinical brain damage in hypertensive patients.

Patients and methods: 101 patients with essential hypertension and 53 controls with no clinical evidence of neurological disease were recruited. Serum concentrations for two brain specific proteins (S100B and neuron specific enolase, NSE) and for inflammatory markers (C-reactive protein, CRP; α 1-antitrypsin, AAT; C3 and C4 complements) were determined. Target organ damage (TOD) to the brain, heart and kidneys was evaluated in HT patients. Patient and Institutional Review Board approval were obtained.

Results: Multiple regression analysis revealed that only NSE and CRP were independently associated with the condition of being hypertensive. NSE was associated to diastolic blood pressure and grade of retinopathy. TOD to the 3 organs was evaluated in 34 patients, showing damage to the brain in 70.6%, heart in 58.5% and kidneys in 50%. NSE was associated with more severe MRI damage (white matter hyperintensities), while inflammatory markers were not related to TOD. After ROC analysis, NSE (cutoff level: 11 μ g/L) was found to predict more severe MRI lesions with 84% sensitivity and 55% specificity.

Conclusion: Serum NSE could constitute a starting point for future investigations in the field of blood based biomarkers for the detection of subclinical brain damage in arterial hypertension.

doi:10.1016/j.jns.2015.08.341

266
WFN15-0726
Stroke 3
Imaging assessment of post cardiac arrest hypoxic ischemic encephalopathy

M. Bertoni^a, M. Snazelle^b, D. Pandit^b, T. Webb^c, R. Dulai^b, Z. Aguirre-Arrizubieta^d. ^aDepartment of Radiology, East Kent Hospitals University NHS Foundation Trust, Ashford, United Kingdom; ^bIntensive Care Unit, East

Kent Hospitals University NHS Foundation Trust, Ashford, United Kingdom; ^cDepartment of Stroke Medicine, East Kent Hospitals University NHS Foundation Trust, Ashford, United Kingdom; ^dDepartment of Electrophysiology, East Kent Hospitals University NHS Foundation Trust, Ashford, United Kingdom

Purpose: To evaluate imaging changes in subjects with hypoxic ischemic encephalopathy (HIE) post cardiac arrest.



Method: From 2013 to 2014, 6 subjects admitted to Intensive Care Unit after successful post cardiac arrest cardiopulmonary resuscitation, were assessed with imaging which included baseline plain computed tomography (CT) and also with perfusion CT (pCT) and magnetic resonance imaging MRI, given lack of improvement of their Glasgow coma scores (GCS) after induced hypothermia. They were also assessed clinically, neurologically and with electroencephalography.

Cortical flow was assessed on pCT with relative blood flow (rBF), relative blood volume (rBV), time to peak (TTP) and mean transit time (MTT) series, following the same protocol in use to assess acute stroke subjects at our institution. MRI included diffusion weighted imaging (DWI) series and apparent diffusion coefficient (ADC) maps. Imaging findings were correlated with clinical assessment. There were no control subjects given the particular clinical situation triggering the protocol mentioned above.

Results: All subjects showed ischemic-type signal intensity changes in cortical areas of the posterior circulation and basal ganglia, which matched an abnormally increased rBFV showing flow rates above 80 ml/min/100 g, when scans were obtained in a period of 48 hours

post induced hypothermia. One of them had a follow up pCT 72 hours post induced hypothermia showing a marked generalized flow drop to 22 ml/min/100 g in cortical regions and basal ganglia.

Conclusion: pCT was a reliable indicator of abnormal flow post induced hypothermia in this cohort of subjects not regaining alertness or improving GCS, showing abnormal flow increase post treatment apparently due to impaired self regulation.



doi:10.1016/j.jns.2015.08.342

267

WFN15-0513

Stroke 3

Endovascular treatment of acute middle cerebral artery occlusion – comparison of treatment methods and identification of outcome predictors

R. Herzig^a, K. Blejcharova^b, E. Vitkova^c, M. Roubec^d, D. Sanak^e, A. Tomek^{f,g}, A. Krajina^h, V. Prochazkaⁱ, M. Kocher^j, F. Charvat^k, D. Krajickova^c, M. Kuliha^d, J. Zapletalova^l, M. Valis^c, D. Skoloudik^d. ^aDepartment of Neurosurgery and Neurooncology Comprehensive Stroke Center, Military University Hospital, Prague, Czech Republic; ^bDepartment of Neurology, Vitkovice Hospital, Ostrava, Czech Republic; ^cDepartment of Neurology Comprehensive Stroke Center, Charles University and University Hospital, Hradec Kralove, Czech Republic; ^dDepartment of Neurology Comprehensive Stroke Center, Ostrava University and University Hospital, Ostrava, Czech Republic; ^eDepartment of Neurology Comprehensive Stroke Center, Palacky University and University Hospital, Olomouc, Czech Republic; ^fDepartment of Neurology Stroke Center, Charles University, Prague, Czech Republic; ^gMotol University Hospital, Prague, Czech Republic; ^hDepartment of Radiology Comprehensive Stroke Center, Charles University and University Hospital, Hradec Kralove, Czech Republic; ⁱDepartment of Radiology Comprehensive Stroke Center, Ostrava University and University Hospital, Ostrava, Czech Republic; ^jDepartment of Radiology Comprehensive Stroke