

WITHDRAWN: Estimated glomerular filtration rate predicts incident stroke among ghanaians with diabetes and hypertension

Fred Stephen Sarfo^{a,b,*}, Linda Meta Mobula^{c,d}, Osei Sarfo-Kantanka^{a,b}, Sheila Adamu^{a,b}, Jacob Plange-Rhule^e, Daniel Ansong^{a,b}, Rexford Adu Gyamfi^f, James Duah^g, Bertha Abraham^h, David Ofori-Adjeiⁱ

^a*Department of Medicine, Kwame Nkrumah University of Science & Technology, Kumasi, Ghana*

^b*Department of Medicine, Komfo Anokye Teaching Hospital, Kumasi, Ghana*

^c*Johns Hopkins University School of Medicine, Baltimore, MD, USA*

^d*Johns Hopkins University Bloomberg School of Public Health, Baltimore, MD, USA*

^e*Ghana College of Physicians and Surgeons, Accra, Ghana*

^f*Agogo Presbyterian Hospital, Agogo, Ghana*

^g*Kings Medical Center, Botanga, Ghana*

^h*Atua Government Hospital, Somanya, Ghana*

ⁱ*Department of Medicine & Therapeutics, University of Ghana, School of Medicine and Dentistry, Accra, Ghana*

This article has been withdrawn at the request of the editor. The Publisher apologizes for any inconvenience this may cause.

The full Elsevier Policy on Article Withdrawal can be found at <https://www.elsevier.com/about/our-business/policies/article-withdrawal>